

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61F 9/007	A1	(11) International Publication Number: WO 00/64391
		(43) International Publication Date: 2 November 2000 (02.11.00)

(21) International Application Number: PCT/US00/11215

(22) International Filing Date: 26 April 2000 (26.04.00)

(30) Priority Data:
60/131,030 26 April 1999 (26.04.99) US

(71)(72) Applicants and Inventors: LYNCH, Mary, G. [US/US];
3845 Club Drive, NE, Atlanta, GA 30319 (US). BROWN,
Reay, H. [US/US]; 3845 Club Drive, NE, Atlanta, GA 30319
(US).

(74) Agents: WARREN, William, L. et al.; Sutherland Asbill &
Brennan LLP, 999 Peachtree Street, NE, Atlanta, GA 30309
(US).


(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB,
BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM,
DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,
UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT,
LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI,
CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

*Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: STENT DEVICE AND METHOD FOR TREATING GLAUCOMA



(57) Abstract

Stent devices and a method for continuously facilitating the flow of aqueous humor through Schlemm's canal where post-operative patency can be maintained with an indwelling stent device. The stent devices provide uni- or bi-directional flow of aqueous humor within and through Schlemm's canal.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

STENT DEVICE AND METHOD FOR TREATING GLAUCOMA

10 Cross - Reference to Related Applications

This application claims the benefit of U.S. Provisional Application No. 60/131,030, filed April 26, 1999.

15 Technical Field

The present invention is generally directed to a surgical treatment for glaucoma, and relates more particularly to a device and method for continuously maintaining the patency of Schlemm's canal with a trough-like, indwelling stent which can be surgically placed to traverse at least a portion of the circumference of the canal and to facilitate the drainage of aqueous humor therethrough.

Background of the Invention

25

Glaucoma is a significant public health problem, because glaucoma is a major cause of blindness. The blindness that results from glaucoma involves both central and peripheral vision and has a major impact on an individual's ability to lead an independent life.

30

Glaucoma is an optic neuropathy (a disorder of the optic nerve) that usually occurs in the setting of an elevated intraocular pressure. The pressure within the eye increases and this is associated with changes in the appearance ("cupping") and function ("blind spots" in the visual field) of the optic nerve. If the pressure remains high enough for a long enough period of time, total vision loss occurs. High pressure develops in an eye because of an internal fluid imbalance.

The eye is a hollow structure that contains a clear fluid called "aqueous humor." Aqueous humor is formed in the posterior chamber of the eye by the ciliary body at a rate of about 2.5 microliters per minute. The fluid, which is made at a fairly constant rate, then passes around the lens, through the pupillary opening in the iris and into the anterior chamber of the eye. Once in the anterior chamber, the fluid drains out of the eye through two different routes. In the "uveoscleral" route, the fluid percolates between muscle fibers of the ciliary body. This route accounts for approximately ten percent of the aqueous outflow in humans. The primary pathway for aqueous outflow in humans is through the "canalicular" route that involves the trabecular meshwork and Schlemm's canal.

The trabecular meshwork and Schlemm's canal are located at the junction between the iris and the sclera. This junction or corner is called "the angle." The trabecular meshwork is a wedge-shaped structure that runs around the circumference of the eye. It is composed of collagen beams arranged in a three-dimensional sieve-like structure. The beams are lined with a monolayer of cells called trabecular cells. The spaces between the collagen beams are filled with an extracellular substance that is produced by the trabecular cells. These cells also produce enzymes that

degrade the extracellular material. Schlemm's canal is adjacent to the trabecular meshwork. The outer wall of the trabecular meshwork coincides with the inner wall of Schlemm's canal. Schlemm's canal is a tube-like structure that runs around the circumference of the cornea. In
5 human adults, Schlemm's Canal is believed to be divided by septa into a series of autonomous, dead-end canals.

The aqueous fluid travels through the spaces between the trabecular beams, across the inner wall of Schlemm's canal into the canal,
10 through a series of about 25 collecting channels that drain from Schlemm's canal and into the episcleral venous system. In a normal situation, aqueous production is equal to aqueous outflow and intraocular pressure remains fairly constant in the 15 to 21 mmHg range. In glaucoma, the resistance through the canalicular outflow system is
15 abnormally high.

In primary open angle glaucoma, which is the most common form of glaucoma, the abnormal resistance is believed to be along the outer aspect of trabecular meshwork and the inner wall of Schlemm's
20 canal. It is believed that an abnormal metabolism of the trabecular cells leads to an excessive build up of extracellular materials or a build up of abnormally "stiff" materials in this area. Histopathology of glaucoma eyes also demonstrates a collapse of Schlemm's canal. Primary open angle glaucoma accounts for approximately eighty-five percent of all
25 glaucoma. Other forms of glaucoma (such as angle closure glaucoma and secondary glaucomas) also involve decreased outflow through the canalicular pathway but the increased resistance is from other causes such as mechanical blockage, inflammatory debris, cellular blockage, etc.

With the increased resistance, the aqueous fluid builds up because it cannot exit fast enough. As the fluid builds up, the intraocular pressure (IOP) within the eye increases. The increased IOP compresses the axons in the optic nerve and also may compromise the vascular supply to the optic nerve. The optic nerve carries vision from the eye to the brain. Some optic nerves seem more susceptible to IOP than other eyes. While research is investigating ways to protect the nerve from an elevated pressure, the only therapeutic approach currently available in glaucoma is to reduce the intraocular pressure.

10

The clinical treatment of glaucoma is approached in a step-wise fashion. Medication often is the first treatment option. Administered either topically or orally, these medications work to either reduce aqueous production or they act to increase outflow. Currently available medications have many serious side effects including: congestive heart failure, respiratory distress, hypertension, depression, renal stones, aplastic anemia, sexual dysfunction and death. Compliance with medication is a major problem, with estimates that over half of glaucoma patients do not follow their correct dosing schedules.

When medication fails to adequately reduce the pressure, laser trabeculoplasty often is performed. In laser trabeculoplasty, thermal energy from a laser is applied to a number of noncontiguous spots in the trabecular meshwork. It is believed that the laser energy stimulates the metabolism of the trabecular cells in some way, and changes the extracellular material in the trabecular meshwork. In approximately eighty percent of patients, aqueous outflow is enhanced and IOP decreases. However, the effect often is not long lasting and fifty percent

of patients develop an elevated pressure within five years. The laser surgery is not usually repeatable. In addition, laser trabeculoplasty is not an effective treatment for primary open angle glaucoma in patients less than fifty years of age, nor is it effective for angle closure glaucoma and
5 many secondary glaucomas.

If laser trabeculoplasty does not reduce the pressure enough, then filtering surgery is performed. With filtering surgery, a hole is made in the sclera and angle region. This hole allows the aqueous fluid to leave
10 the eye through an alternate route.

The most commonly performed filtering procedure is a trabeculectomy. In a trabeculectomy, a posterior incision is made in the conjunctiva, the transparent tissue that covers the sclera. The conjunctiva
15 is rolled forward, exposing the sclera at the limbus. A partial thickness scleral flap is made and dissected half-thickness into the cornea. The anterior chamber is entered beneath the scleral flap and a section of deep sclera and trabecular meshwork is excised. The scleral flap is loosely sewn back into place. The conjunctival incision is tightly closed.
20 Post-operatively, the aqueous fluid passes through the hole, beneath the scleral flap and collects in an elevated space beneath the conjunctiva. The fluid then is either absorbed through blood vessels in the conjunctiva or traverses across the conjunctiva into the tear film.

25 Trabeculectomy is associated with many problems. Fibroblasts that are present in the episclera proliferate and migrate and can scar down the scleral flap. Failure from scarring may occur, particularly in children and young adults. Of eyes that have an initially successful trabeculectomy, eighty percent will fail from scarring within

three to five years after surgery. To minimize fibrosis, surgeons now are applying antifibrotic agents such as mitomycin C (MMC) and 5-fluorouracil (5-FU) to the scleral flap at the time of surgery. The use of these agents has increased the success rate of trabeculectomy but also has
5 increased the prevalence of hypotony. Hypotony is a problem that develops when aqueous flows out of the eye too fast. The eye pressure drops too low (usually less than 6.0 mmHg); the structure of the eye collapses and vision decreases.

10 Trabeculectomy creates a pathway for aqueous fluid to escape to the surface of the eye. At the same time, it creates a pathway for bacteria that normally live on the surface of the eye and eyelids to get into the eye. If this happens, an internal eye infection can occur called endophthalmitis. Endophthalmitis often leads to permanent and profound
15 visual loss. Endophthalmitis can occur anytime after trabeculectomy. The risk increases with the thin blebs that develop after MMC and 5-FU. Another factor that contributes to infection is the placement of a bleb. Eyes that have trabeculectomy performed inferiorly have about five times the risk of eye infection than eyes that have a superior bleb. Therefore,
20 initial trabeculectomy is performed superiorly under the eyelid, in either the nasal or temporal quadrant.

In addition to scarring, hypotony and infection, there are other complications of trabeculectomy. The bleb can tear and lead to
25 profound hypotony. The bleb can be irritating and can disrupt the normal tear film, leading to blurred vision. Patients with blebs generally cannot wear contact lenses. All of the complications from trabeculectomy stem from the fact that fluid is being diverted from inside the eye to the external surface of the eye.

When trabeculectomy doesn't successfully lower the eye pressure, the next surgical step often is an aqueous shunt device. An aqueous shunt device of the prior art is a silicone tube that is attached at one end to a plastic (polypropylene or other synthetic) plate. With an aqueous shunt device, an incision is made in the conjunctiva, exposing the sclera. The plastic plate is sewn to the surface of the eye posteriorly, usually over the equator. A full thickness hole is made into the eye at the limbus, usually with a needle. The tube is inserted into the eye through this hole. The external portion of the tube is covered with either donor sclera or pericardium. The conjunctiva is replaced and the incision is closed tightly. Many problems exist with the current technology of aqueous shunt devices including scarring, failure, hypotony, and infection.

Some prior art references for glaucoma management have been directed at Schlemm's canal, but these have not involved the placement of long-term, indwelling stents. For example, U.S. Patent No. 5,360,399 teaches the placement of a portion of a plastic or steel tube in Schlemm's canal with injection of a viscous material through the tube to hydraulically dissect the trabecular meshwork. The tube is removed from the canal following injection. Furthermore, relative to that portion within Schlemm's canal, the '399 device has a larger diameter injection cuff element, which serves as an adapter for injection and irrigation. Therefore, this device is not adapted for permanent placement within Schlemm's canal.

A need exists, then, for a more physiologic system to enhance the drainage of aqueous fluid through Schlemm's canal.

Enhancing aqueous flow directly into Schlemm's canal would minimize scarring since the angle region is populated with a single line of nonproliferating trabecular cells. Enhancing aqueous flow directly into Schlemm's canal would minimize hypotony since the canal is part of the normal outflow system and is biologically engineered to handle the normal volume of aqueous humor. Enhancing aqueous flow directly into Schlemm's canal would eliminate complications such as endophthalmitis and leaks.

10 **Summary of the Invention**

The present invention is directed to a novel stent and an associated surgical method for the surgical correction of glaucoma in which the stent is placed into Schlemm's canal to expand the canal's dimensions and maintain its patency. The present invention therefore facilitates the normal physiologic pathway for drainage of aqueous humor into and through Schlemm's canal. The present invention is further directed to providing a permanent, indwelling stent within Schlemm's canal for glaucoma management.

20

Brief Description of the Drawings

FIG. 1 is an illustration showing a side view of one embodiment of the present invention, in which the inventive stent is comprised of tubular elements traversing the circumference of Schlemm's canal.

FIG. 2 is an illustration showing another embodiment of the present invention, in which the inventive stent is comprised of luminal mesh tubular elements.

5 FIG. 3 is an illustration showing another embodiment of the present invention in which the inventive stent is comprised of elements that are partially tubular and partially open in their configuration.

10 FIG. 4 is an illustration showing the anatomic details of the human eye.

15 FIG. 5 is an illustration showing the anatomic relationships (not to scale) of the surgical placement of an exemplary embodiment of the present invention.

Detailed Description of Present Invention

20 The present invention provides an aqueous humor stent device to be placed within a portion of Schlemm's canal of the eye as an indwelling implant to expand and maintain the patency of the canal, in which the stent device comprises a body portion shaped to be wholly received within Schlemm's canal to facilitate the natural drainage of aqueous humor to the collecting channels of the eye.

25

 The present invention also provides embodiments of an inventive stent comprising a thin body of biocompatible material of a length and shape adapted to be wholly received within Schlemm's canal and to extend within a portion of the circumference of Schlemm's canal,

and having a channel therein to facilitate the passage of aqueous humor into and through Schlemm's canal to the collecting channels. The invention contemplates many different configurations for a stent device, provided that each assists in channeling aqueous humor throughout Schlemm's canal, such as by providing a lumen, trough, wick or capillary action. In some embodiments of the invention, the body of the stent can move between a first insertion position and a second expanded stenting position when in a desired location of the canal.

10 The present invention also provides methods of use of the stent devices. One embodiment of the present invention is directed to a surgical method to implant the inventive stent into a portion of the circumference of Schlemm's canal. The device extending into Schlemm's canal may be fashioned from a flexible, porous or non-
15 porous, biologically inert material to approximately equal a portion of the radius, curvature, and diameter of Schlemm's canal. All or parts of the device may be tubular or non-tubular, and fenestrated or non-fenestrated. The device may further be sized to allow placement through all or some of the circumference of Schlemm's canal.

20 Traditional glaucoma teaching states that Schlemm's canal in an adult is divided by septa into separate canals, rendering the complete passage of a suture impossible. Preliminary studies on adult human eye bank eyes have shown that Schlemm's canal is, indeed, patent. A suture
25 can be passed through the entire circumference of the canal. It has not been heretofore determined that Schlemm's canal is patent throughout its circumference in normal individuals, as opposed to being divided by septa into multiple dead end canals. The present invention utilizes this

knowledge to create and maintain patency within Schlemm's canal with the present stent devices.

One embodiment of the present invention is illustrated in
5 **FIG. 1**, in which the stent device **100** is shown in a side view. The stent device **100** is comprised of a tubular body portion **10** defining a lumen **5** which may have solid tubular walls or may contain a plurality of fenestrations **15** communicating between the lumen **5** and the exterior. The body portion has a pre-formed curvature with a radius r which
10 approximates the 6 mm radius of Schlemm's canal of an adult human eye. The cross-sectional diameter of the body portion **10** is sized to be fully received within Schlemm's canal. The body portion **10** may be either an enclosed tubular or multisided structure, or it may be a flat, angular, or curved open structure, or some combination of the above
15 when cross-sectioned at different sites along the entirety of its length. The fenestrations **15** may be placed along any portion of the device **100** to facilitate the passage of fluid therethrough.

Other examples of embodiments of the present invention are
20 shown in **FIGS. 2-3**. **FIG. 2** shows an embodiment of the inventive stent in which the device **100** comprises a luminal tubular mesh in its configuration, again with a pre-formed curvature with a radius r to approximate the 6 mm radius of Schlemm's canal and a cross-sectional diameter of the body portion **10** sized to be fully received within
25 Schlemm's canal.

FIG. 3 shows an embodiment of the inventive stent in which the body portion **10** is open and curved throughout its length in a trough-like channel, again with a pre-formed curvature with a radius r to

approximate the 6 mm radius of Schlemm's canal and a cross-sectional diameter of the body portion 10 sized to be fully received within Schlemm's canal.

5 As the inventive device is a long-term implant, it can be fabricated from a material that will be innocuous to the tissues and fluids with which it is in contact. It is preferable that the device not be absorbed, corroded, or otherwise structurally compromised during its *in situ* tenure. Moreover, it is equally preferable that the eye tissues and the
10 aqueous remain non-detrimentally affected by the presence of the implanted device. A number of materials are available to meet the engineering and medical specifications for the stents. In the exemplary embodiments of the present invention, the stent device 100 is constructed of a biologically inert, flexible material such as silicone or similar
15 polymers. Alternate materials might include, but are not limited to, thin-walled polytetrafluoroethylene, polypropylene or other polymers. Other metals and alloys known in the art of stenting can also be used, such as stainless steel, titanium or nitinol. The stent can also be fabricated with therapeutic agents that migrate from the device over time.

20

 In the embodiments shown in FIGS. 1-5, the body portion 10 may have a pre-formed curve to approximate the 6.0 mm radius of Schlemm's canal in a human eye. The body portion 10 may be of sufficient length to extend through any length of the entire circumference
25 of Schlemm's canal, with a total length for the body portion 10 of about 1.0 mm to 40 mm, or about 2 mm to 20 mm, or about 5 mm to permit circumferential placement through Schlemm's canal. The diameter or width of the body portion 10 can be sized to yield an internal diameter of between 0.1 mm and 0.5 mm, preferably about 0.2 mm and an external

diameter of between 0.1 mm and 0.5 mm, or about 0.3 mm, for a tubular or curved stent, or a comparable maximal width for a stent with a multiangular configuration. The body portion 10 may contain a plurality of fenestrations to allow fluid egress, arranged to prevent occlusion by the adjacent walls of Schlemm's canal, particularly in the direction of the collecting channels.

The surgical anatomy relevant to the present invention is illustrated in FIG. 4. Generally, FIG. 4 shows the anterior chamber 35, Schlemm's canal 30, the iris 40, cornea 45, trabecular meshwork 50, collecting channels 55, episcleral veins 60, pupil 65, and lens 70. FIG. 5 illustrates the surgical placement of the exemplary embodiment of the present invention, with the relevant anatomic relationships. It should be noted that the inventive device is designed so that placement of multiple stents within Schlemm's canal 30 can result in a near-circumferential traverse of Schlemm's canal 30. The surgical incision into Schlemm's canal 30 is closed, with no direct external communication with the stent device 100 intended.

The surgical procedure necessary to insert the device may include all or some of the following steps: A conjunctival incision is made. A partial thickness scleral flap is then created and dissected half-thickness into clear cornea. The posterior aspect of Schlemm's canal is identified and the canal is entered posteriorly. The anterior chamber may be deepened with injection of a viscoelastic or miotic agent. A balloon catheter, such as described in U.S. Serial No. _____, filed April 26, 2000, may be introduced into Schlemm's canal, and inflated to dilate portions of Schlemm's canal, followed by the selective deflation of the balloon and placement of one or more stent devices into Schlemm's canal.

Alternately, the stent devices may be introduced directly on a balloon catheter device. Therefore, a plurality of stent segments may be placed at selected locations along the circumference of Schlemm's canal. Any residual stent material is trimmed, and the scleral flap and conjunctival
5 wound are closed in a conventional manner.

While the above-described embodiments are exemplary, the invention contemplates a wide variety of shapes and configurations of the stent to provide fluid communication between the anterior chamber and
10 Schlemm's canal and to the collecting channels. The above-described embodiments are therefore not intended to be limiting to the scope of the claims and equivalents thereof.

CLAIMS

What is claimed is:

1. A stent device to expand and maintain the patency of Schlemm's canal in the eye, the stent device comprising a body portion shaped to be wholly received within a portion of Schlemm's canal, wherein said stent device improves fluid communication between the anterior chamber and Schlemm's canal.
2. The stent device of Claim 1, wherein the body portion of the stent has a curve having a radius which approximates the radius of Schlemm's canal of a human eye.
3. The stent device of Claim 2, wherein said curve has a radius of between about 3 mm and 10 mm.
4. The stent device of Claim 2, wherein said curve has a radius of about 6 mm.
5. The surgical device of Claim 1, wherein said body portion is at least partially luminal having an outer diameter of about 0.1 mm to 0.5 mm.
6. The stent device of Claim 1, wherein said body portion is at least partially luminal having an outer diameter of about 0.3 mm.
7. The stent device of Claim 1, wherein said body portion has a length of about 1 mm to 40 mm.
8. The stent device of Claim 1, wherein said body portion has a length of about 20 mm.

9. The stent device of Claim 1, where the body portion of the stent has a plurality of fenestrations therein that allow the passage of fluid into Schlemm's canal.
10. The stent device of Claim 1, wherein the body is curved to define a trough-like, partially open channel along at least some of the length of said body open toward the collecting channels of the eye.
11. A stent device for use in the eye to relieve excess intraocular pressure by facilitating drainage through and within Schlemm's canal, comprising:
a thin body of biocompatible material of a length and shape adapted to be wholly retained within a portion of Schlemm's canal and having a channel therein to facilitate the passage of aqueous humor therealong.
12. The stent device of Claim 11, wherein the body portion provides for drainage in both directions along Schlemm's canal.
13. The stent device of Claim 11, wherein the body portion of the stent has a curve having a radius which approximates the radius of Schlemm's canal of a human eye.
14. The stent device of Claim 13, wherein said curve has a radius of between about 3 mm and 10 mm.
15. The stent device of Claim 14, wherein said curve has a radius of about 6 mm.
16. The stent device of Claim 11, wherein said body portion has an outer diameter of about 0.1 mm to 0.5 mm.

17. The stent device of Claim 11, wherein said body portion has an outer diameter of about 0.3 mm.
18. The stent device of Claim 11, wherein said body portion has a length of about 1 mm to 40 mm.
19. The stent device of Claim 11, wherein said body portion has a length of about 20 mm.
20. The stent device of Claim 11, where the body portion of the stent has a plurality of fenestrations therein that allow the passage of fluid into Schlemm's canal.
21. The stent device of Claim 11, wherein the body is curved to define a trough-like, partially open channel along at least some of the length of said body open toward the collecting channels.
22. A method for the surgical treatment of glaucoma, comprising, performing a trabeculotomy through a conjunctival flap made at the limbus; developing a partial thickness scleral flap; radially incising the junction between the angle tissue and the sclera, which is surgically extended until Schlemm's canal is entered posteriorly, and placing one or more of the stent devices of Claim 1 within Schlemm's canal.
23. A method for the surgical treatment of glaucoma, comprising, performing a trabeculotomy through a conjunctival flap made at the limbus; developing a partial thickness scleral flap; radially incising the junction between the angle tissue and the sclera, which is surgically extended until Schlemm's canal is

entered posteriorly, and placing one or more of the stent devices of Claim 11 within Schlemm's canal.

1/3




Fig. 1




Fig. 2

2/3




Fig. 3




Fig. 5




Fig. 4

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/11215

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61F9/007

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 372 577 A (UNGERLEIDER BRUCE A) 13 December 1994 (1994-12-13) figure 9 column 4, line 47 - line 68	1, 11
A	US 5 073 163 A (LIPPMAN MYRON E) 17 December 1991 (1991-12-17) figure 3 column 3, line 3 - line 55 claim 1	1, 11
A	US 4 750 901 A (MOLTENO ANTHONY C B) 14 June 1988 (1988-06-14) figures 1-3 claims 1-7	1, 11
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

8 September 2000

Date of mailing of the international search report

18/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 851 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Mary, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/11215

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 037 604 A (NEWKIRK JOHN B) 26 July 1977 (1977-07-26) figure 5 column 2, line 6 -column 3, line 14	1,11
A	US 3 788 327 A (DONOWITZ H) 29 January 1974 (1974-01-29) column 2, line 29 -column 3, line 2 figures 1-4	1,11
P,A	DE 198 40 047 A (NEUHANN THOMAS) 23 March 2000 (2000-03-23) figures 1,2 column 4, line 33 -column 5, line 11	1,11

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/11215

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5372577 A	13-12-1994	US 4936825 A	26-06-1990
US 5073163 A	17-12-1991	NONE	
US 4750901 A	14-06-1988	NZ 215409 A	24-02-1989
		GB 2187963 A, B	23-09-1987
US 4037604 A	26-07-1977	NONE	
US 3788327 A	29-01-1974	NONE	
DE 19840047 A	23-03-2000	WO 0013627 A	16-03-2000